COMPOSITIONAL REGRESSION-BASED METHODS FOR SST RECONSTRUCTION



Alpine and Mediterranean Quaternary, 33 (1), 2020, 31-41

https://doi.org/10.26382/AMQ.2020.02

ISSN 2279-7335

Supplementary Appendix

Supplementary Table

method	R ² (squared r- coefficient)	RMSE (°C)	RMSD (°C)
CoDa-PLSR			
15 taxa 2 groups annual	0.9351 (0.9718)	1.3349	
15 taxa 2 groups robust annual	0.9060 (0.9591)	1.7571	
15 taxa 2 groups summer	0.8796 (0.9533)	1.6567	2.3007
15 taxa 2 groups winter	0.9206 (0.9609)	1.5964	
19 taxa 2 groups annual	0.9315 (0.9702)	1.3716	
19 taxa 2 groups summer	0.8741 (0.9512)	1.6945	2.3414
19 taxa 2 groups winter	0.9186 (0.9599)	1.6158	
15 taxa annual	0.9367	1.9985	
15 taxa summer	0.917	2.2054	3.1871
15 taxa winter	0.9188	2.3008	
19 taxa annual	0.9326	2.0625	
CoDa-PCR			
15 taxa 2 groups annual	0.9276 (0.9686)	1.5651	
15 taxa 2 groups summer	0.8678 (0.9487)	1.7364	2.3927
15 taxa 2 groups winter	0.8781 (0.9645)	1.6462	
15 taxa 2 groups robust	0.9262 (0.9685)	1.4103	
15 taxa annual	0.9210	2.2323	
15 taxa summer	0.8933	2.5071	3.4350
15 taxa winter	0.9149	2.3621	
Coda-MAT			
CoDa-MAT annual	0.9857	0.9484	

Cross-validation results for ilr/clr-PLSR ilr/clr-PCR, obtained under different conditions.