



## Research Paper

# Permanent cystathione- $\beta$ -Synthase gene knockdown promotes inflammation and oxidative stress in immortalized human adipose-derived mesenchymal stem cells, enhancing their adipogenic capacity



Ferran Comas <sup>a</sup>, Jèssica Latorre <sup>a</sup>, Francisco Ortega <sup>a</sup>, Núria Oliveras-Cañellas <sup>a</sup>, Aina Lluch <sup>a</sup>, Wifredo Ricart <sup>a</sup>, José Manuel Fernández-Real <sup>a,b,\*\*</sup>, José María Moreno-Navarrete <sup>a,b,\*</sup>

<sup>a</sup> Department of Diabetes, Endocrinology and Nutrition, Institut d'Investigació Biomèdica de Girona (IdIBGI), CIBEROBN (CB06/03/010) and Instituto de Salud Carlos III (ISCIII), Girona, Spain

<sup>b</sup> Department of Medicine, Universitat de Girona, Girona, Spain

## ARTICLE INFO

**Keywords:**

Human adipose-derived mesenchymal stem cells  
Cystathionine  $\beta$ -synthase  
Cellular senescence  
Inflammation  
Oxidative stress and adipogenesis

## ABSTRACT

In the present study, we aimed to investigate the impact of permanent cystathione- $\beta$ -Synthase (CBS) gene knockdown in human telomerase reverse transcriptase (hTERT) immortalized human adipose-derived mesenchymal stem cells (ASC52telo) and in their capacity to differentiate into adipocytes. CBS gene KD in ASC52telo cells led to increased cellular inflammation (*IL6*, *CXCL8*, *TNF*) and oxidative stress markers (increased intracellular reactive oxygen species and decreased reduced glutathione levels) in parallel to decreased H<sub>2</sub>S production and rejuvenation (*LC3* and *SIRT1*)-related gene expression. In addition, CBS gene KD in ASC52telo cells resulted in altered mitochondrial respiratory function, characterised by decreased basal respiration (specifically proton leak) and spare respiratory capacity, without significant effects on cell viability and proliferation. In this context, shCBS-ASC52telo cells displayed enhanced adipogenic (*FABP4*, *ADIPOQ*, *SLC2A4*, *CEBPA*, *PPARG*), lipogenic (*FASN*, *DGAT1*)- and adipocyte (*LEP*, *LBP*)-related gene expression markers, decreased expression of proinflammatory cytokines, and increased intracellular lipid accumulation during adipocyte differentiation compared to control ASC52telo cells. Otherwise, the increased adipogenic potential of shCBS-ASC52telo cells was detrimental to the ability to differentiate into osteogenic lineage. In conclusion, this study demonstrated that permanent CBS gene KD in ASC52telo cells promotes a cellular senescence phenotype with a very increased adipogenic potential, promoting a non-physiological enhanced adipocyte differentiation with excessive lipid storage.

## 1. Introduction

Recent studies in 3T3-L1 cells pointed to a relevant role of transulfuration pathway in adipogenesis [1–4]. Even though Tsai et al. reported that both transulfuration pathway enzymes [cystathionine  $\beta$ -synthase (CBS) and cystathionine  $\gamma$ -lyase (CTH or CSE)] were important in adipocyte differentiation [1], the latter studies demonstrated that only CTH exerted a relevant role in adipogenesis increasing PPAR $\gamma$  activity during adipocyte differentiation [2,3]. In fact, in a previous study, in which *Cth* and *Cbs* gene expression in 3T3-L1 cells was analysed and

compared, we found that *Cth* was highly expressed in preadipocytes and increased during adipocyte differentiation, whereas expression of *Cbs* gene was almost undetectable in preadipocytes and did not change during adipocyte differentiation, even tending to decrease in the first two days of the process [4].

Human adipose-derived mesenchymal stem cells (hAMSC), the adipocyte precursor cells located in adipose tissue stromal vascular fraction, are required for adipose tissue hyperplasia and functionality [5, 6]. A previous study reported a useful human telomerase reverse transcriptase (hTERT) immortalized hAMSC (ASC52telo) cell model to study

\* Corresponding author. Department of Diabetes, Endocrinology and Nutrition, Institut d'Investigació Biomèdica de Girona (IdIBGI), Parc Hospitalari Martí i Julià - Edifici M2, C/ Dr. Castany s/n, 17190, Salt, Spain.

\*\* Corresponding author. Section of Diabetes, Endocrinology and Nutrition, Institut d'Investigació Biomèdica de Girona (IdIBGI), Hospital of Girona "Dr Josep Trueta", Carretera de França s/n, 17007, Girona, Spain.

E-mail addresses: [jmfreal@idibgi.org](mailto:jmfreal@idibgi.org) (J.M. Fernández-Real), [jmoreno@idibgi.org](mailto:jmoreno@idibgi.org) (J.M. Moreno-Navarrete).


<https://doi.org/10.1016/j.redox.2020.101668>

Received 23 June 2020; Accepted 29 July 2020

Available online 2 August 2020

2213-2317/© 2020 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



**Fig. 1.** A) CBS and CTH mRNA levels in ASC52telo cells, \*\*\*p < 0.001. B) Representative Western blot of CBS and CTH protein in ASC52telo cell lysates. C-F) ADIPOQ, SLC2A4, CTH and CBS mRNA levels during adipocyte differentiation of ASC52telo cells. \*p < 0.05 and \*\*p < 0.01 vs day 0.

adipocyte differentiation [7].

To the best of our knowledge, the importance of transulfuration pathway in hAMSC adipogenesis has not been yet examined. Taking advantage of ASC52telo cells, in the present study, we aimed to investigate the possible role of CBS enzyme in hAMSC during adipocyte differentiation.

## 2. Material and methods

### 2.1. Differentiation of human immortalized adipose-derived mesenchymal stem cells

Human telomerase reverse transcriptase immortalized adipose-derived MSC (ASC52telo, SCRC-4000, ATCC, LGC Standards SLU, Barcelona, Spain) cells were cultured in Mesenchymal Stem Cell Basal Medium (ATCC PCS-500-030) plus FBS (2%), rhFGF basic (5 ng/ml), rhFGF acidic (5 ng/ml), rhEGF (5 ng/ml), L-Alanyl-L-Glutamine (2.4 mM) and G418 (0.2 mg/ml) at 37 °C in a 5% CO<sub>2</sub> in air atmosphere.

**Adipogenic differentiation.** ASC52telo cells were cultured in three repetitive cycles of 72 h in adipogenic differentiation medium composed of DMEM/Nutrient Mix F-12 medium, FBS (10%), penicillin, streptomycin, human insulin (10 µg/mL), DXM (1 µmol/L), isobutylmethylxanthine (0.5 mmol/L), and PPAR $\gamma$  agonists (rosiglitazone, 1 µmol/L), followed by 72 h in adipogenic maintenance medium composed of DMEM/Nutrient Mix F-12 medium, FBS (10%), penicillin, streptomycin, and human insulin (10 µg/mL).

**Osteogenic differentiation.** ASC52telo cells were cultured with osteogenic differentiation medium composed of DMEM/Nutrient Mix F-12 medium, FBS (10%), 50 µg/ml ascorbic 2-phosphate, 10 nM dexamethasone and 10 mM  $\beta$ -glycerophosphate. Osteogenic medium was refreshed every 3 days. After 9 days, the cells were harvested.

### 2.2. Short hairpin (sh) RNA-mediated knockdown of CBS gene

Permanent silencing was performed using CBS-targeted and control (scrambled) shRNA lentiviral particles (sc-60335-V and sc-108080,

Santa Cruz Biotechnology, CA, USA) and following the manufacturer instructions. Positive hAMSC harboring shRNA cassette for CBS were selected by puromycin (3 µg/mL) selection 60 h after infection.

### 2.3. In vitro measurements


Intracellular reactive oxygen species were measured using Fluorometric Intracellular ROS Kit (Cat. n° MAK142, Sigma, Madrid, Spain), reduced and total glutathione was measured using Glutathione Colorimetric Assay Kit (Cat. n° K261-100, Biovision, CA, USA). LDH activity and 3-(4,5-dimethyl-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay were performed with specific commercial kits, Cytotoxicity Detection Kit (LDH) (Cat. n° 11644793001, Roche Diagnostics SL, Barcelona, Spain) and Cell Proliferation Kit (MTT) (Cat. n° 11465007001, Roche Diagnostics SL, Barcelona, Spain). H<sub>2</sub>S concentration in cultured medium was assessed as detailed elsewhere [4], using a naphthalimide-based fluorescent sensor 6-Azido-2-[2-[2-(2-hydroxyethoxy)ethoxy]ethyl]benzo[de]isoquinoline-1,3-dione (L1), which was chemically synthesised in Institute of Computational Chemistry and Catalysis (Chemistry Department, University of Girona) as described previously [8]. Intracellular lipid accumulation was assessed by Oil Red O staining as detailed elsewhere [4].

### 2.4. Mitochondrial respiratory function

Mitochondrial respiratory function was assessed using Seahorse XFP Extracellular Flux Analyzer (Seahorse Bioscience, Agilent Technologies) using Seahorse XFP Cell Mito Stress Test Kit as detailed elsewhere [9].

### 2.5. RNA expression

Briefly, RNA purification was performed using RNeasy Lipid Tissue Mini Kit (QIAgen, Izasa SA, Barcelona, Spain) and the integrity was checked by Agilent Bioanalyzer (Agilent Technologies, Palo Alto, CA). Gene expression was assessed by real time PCR using an LightCycler® 480 Real-Time PCR System (Roche Diagnostics SL, Barcelona, Spain),



**Fig. 2. A-M)** Effect of permanent CBS gene knockdown (shCBS) on CBS mRNA (A) and CBS protein (B) levels, IL6 (C), CXCL8 (D) and TNF (E) mRNA levels, intracellular ROS levels (F), GSH/total glutathione (G), H<sub>2</sub>S production (H), LC3 (I) and SIRT1 (J) mRNA levels, mitochondrial respiration (K), MTT assay (L) and LDH activity (M) in ASC52telo cells. \*p < 0.05, \*\*p < 0.01 and \*\*\*p < 0.001 vs shC-ASC52telo cells.

using TaqMan® and SYBRgreen technology suitable for relative genetic expression quantification. The commercially available and pre-validated TaqMan® primer/probe sets used were as follows: Peptidylprolyl isomerase A (cyclophilin A) (4333763, PPIA as endogenous control), cystathionine γ-lyase (CTH, Hs00542284\_m1), cystathionine β-synthase (CBS, Hs00163925\_m1), adiponectin (ADIPOQ, Hs00605917\_m1), peroxisome proliferator-activated receptor gamma (PPARG, Hs00234592\_m1), fatty acid synthase (FASN, Hs00188012\_m1), diacylglycerol O-acyltransferase 1 (DGAT1, Hs01020362\_g1), CCAAT/enhancer binding protein alpha (CEBPA, Hs00269972\_s1), solute carrier family 2 member 4 (SLC2A4 or GLUT4, Hs00168966\_m1), insulin receptor substrate 1 (IRS1, Hs00178563\_m1), Leptin (LEP, Hs00174877\_m1), lipopolysaccharide binding protein (LBP, Hs01084621\_m1), interleukin 6 (interferon, beta 2) (IL6, Hs00985639\_m1), tumor necrosis factor (TNF, Hs00174128\_m1), C-X-C motif chemokine ligand 8 (CXCL8 or IL8, Hs00174103\_m1), C-C motif chemokine ligand 2 (CCL2 or MCP1, Hs00234140\_m1), microtubule associated protein 1 light chain 3 alpha (MAP1LC3A or LC3, Hs00738808\_m1), sirtuin 1 (SIRT1, Hs01009005\_m1) and fatty acid binding protein 4, adipocyte (FABP4, Hs01086177\_m1).


#### 2.6. Protein preparation and Western blot analysis

Cellular protein were extracted directly in radioimmuno-

precipitation assay (RIPA) buffer (0.1% SDS, 0.5% sodium deoxycholate, 1% Nonidet P-40, 150 mM NaCl, and 50 mM Tris-HCl, pH 8.0), supplemented with protease inhibitors (1 mM phenylmethylsulfonyl fluoride). Cellular debris and lipids were eliminated by centrifugation of the solubilized samples at 13000 rpm for 10 min at 4 °C, recovering the soluble fraction. Protein concentration was determined using the RC/DC Protein Assay (Bio-Rad Laboratories, Hercules, CA). RIPA protein extracts (25 µg) were separated by SDS-PAGE and transferred to nitrocellulose membranes by conventional procedures. Membranes were immunoblotted with anti-CTH, CBS and β-actin (sc-365382, sc-133154, sc-47778, Santa Cruz Biotechnology, CA, USA). Anti-mouse IgG coupled to horseradish peroxidase was used as a secondary antibody. Horseradish peroxidase activity was detected by chemiluminescence, and quantification of protein expression was performed using scion image software.

#### 2.7. Statistical analyses

Statistical analyses were performed using SPSS 12.0 software. Unpaired t-test and nonparametric test (Mann Whitney test) was used to analyse *in vitro* experimental data. Levels of statistical significance were set at p < 0.05.



**Fig. 3. A-L)** Effect of permanent *CBS* gene knockdown (shCBS) on *CBS*, *FABP4*, *ADIPOQ*, *SLC2A4*, *CEBPA*, *DGAT1*, *FASN*, *IRS1*, *IL6*, *TNF*, *CXCL8* and *CCL2* mRNA levels in a time course experiment during ASC52telo cell adipocyte differentiation. \* $p < 0.05$ , \*\* $p < 0.01$  and \*\*\* $p < 0.001$  vs shC-ASC52telo cells.

### 3. Results

#### 3.1. CBS is much more abundant than CTH in ASC52telo cells

In ASC52telo cells, and contrary to 3T3-L1 cells [4], *Cbs* gene expression and CBS protein levels were significantly increased compared to *CTH* (Fig. 1A and B).

When adipocyte differentiation was examined in ASC52telo cells, we found a significant increase of adipogenic (*ADIPOQ* and *SLC2A4*) gene expression (Fig. 1C and D) in a time course experiment, confirming that adipogenesis has been appropriately induced in these cells. Similar to the observations in 3T3-L1 cells [4], *CTH* gene expression increased during hAMSC adipocyte differentiation in parallel to adipogenic genes (Fig. 1C-E), whereas *CBS* gene expression also increased but only in the last days of adipocyte differentiation process (Fig. 1F).

#### 3.2. CBS gene KD in ASC52telo cells promotes inflammation and oxidative stress


Considering the higher CBS compared CTH protein levels, the effect of permanent *CBS* gene knockdown (KD) on ASC52telo cells was tested. *CBS* gene KD in these cells (Fig. 2A and B) led to a significant increase in cellular inflammation (*IL6*, *CXCL8*, *TNF*) and oxidative stress markers, including increased intracellular reactive oxygen species and decreased intracellular reduced glutathione levels, in parallel to decreased H<sub>2</sub>S production (Fig. 2C-H). In addition, *SIRT1* and *LC3* mRNA were also decreased (Fig. 2I and J). Importantly, when mitochondrial respiration was evaluated, we found that *CBS* gene KD in ASC52telo cells resulted in decreased basal respiration (specifically proton leak) and spare

respiratory capacity (Fig. 2K). Otherwise, no significant effects *CBS* gene knockdown on cell proliferation (MTT assay, Fig. 2L) or necrosis (LDH activity, Fig. 2M) were found.

#### 3.3. CBS gene KD in ASC52telo cells enhances differentiation into adipocytes

In this context, *CBS* gene KD during ASC52telo adipocyte differentiation (Fig. 3A) led to enhanced adipocyte differentiation, with increased expression of adipogenic genes (*FABP4*, *ADIPOQ*, *SLC2A4*, *CEBPA*, *DGAT1*, *FASN*), but decreased *IRS1*, in a time course experiment (Fig. 3B-H; Suppl Figure 1). Consistent with previous findings (Fig. 2C-E), proinflammatory cytokines (*IL6*, *TNF* and *CXCL8*), but not the chemokine *CCL2*, were significantly increased at day 0 (Fig. 3I-L). Of interest, *IL6*, *TNF*, *CXCL8* and *CCL2* mRNA were strongly reduced during adipocyte differentiation, being significantly decreased *IL6* and *CCL2* at day 5 and 9 in shCBS-ASC52telo cells compared to shC-ASC52telo (control) cells (Fig. 3I-L).

To confirm these enhanced adipogenic capacity of shCBS-ASC52telo, in an independent experiment, we evaluated these cells in adipogenic conditions at day 15 (Fig. 4A). Reinforcing previous observations, shCBS-ASC52telo cells displayed enhanced adipogenic (*FABP4*, *ADIPOQ*, *SLC2A4*, *CEBPA*, *PPARG*), lipogenic (*FASN*, *DGAT1*) and adipocyte (*LEP*, *LBP*)-related gene expression markers (Fig. 4B-J), decreased expression of proinflammatory cytokines (*IL6* and *TNF*) (Fig. 4K-L), and increased intracellular lipid accumulation (Fig. 4M-N) at day 15.



**Fig. 4.** A-L) Effect of permanent *CBS* gene knockdown (shCBS) on *CBS*, *FABP4*, *ADIPOQ*, *SLC2A4*, *CEBPA*, *PPARG*, *FASN*, *DGAT1*, *LEP*, *LBP*, *IL6* and *TNF* mRNA levels. M-N) Effect of permanent *CBS* gene knockdown (shCBS) on intracellular lipid accumulation, visually assessed at 10x magnifications (M) and measured by Oil Red O staining (N) at day15 of ASC52telo cell adipocyte differentiation. \* $p < 0.05$ , \*\* $p < 0.01$  and \*\*\* $p < 0.001$  vs shC-ASC52telo cells. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

#### 3.4. *CBS* gene KD in ASC52telo cells decreased osteogenic markers

Next, to evaluate the capacity of shCBS-ASC52telo cells to differentiate into another cell type, these cells were cultured in osteogenic conditions during 9 days. shCBS-ASC52telo cells (Fig. 5A) displayed reduced osteogenic markers (such as *RUNX2*, *BGLAP* and *TFGB1*) (Fig. 5B-D) in parallel to decreased rejuvenation (*LC3* and *SIRT1*)-related genes (Fig. 5E and F), but increased *FABP4* (Fig. 5G) and proinflammatory cytokines (*IL6*, *TNF* and *IL8*) (Fig. 5H-J).


#### 4. Discussion

The current study demonstrated that, contrary to 3T3-L1 cells in which CTH levels were significantly increased but CBS was almost undetectable [4], ASC52telo cells showed an increased CBS/CTH ratio, characterized by increased CBS but very low levels of CTH mRNA and protein. In line with these findings, previous studies also demonstrated increased CBS vs CTH mRNA and protein levels in human mesenchymal stem cells [10,11].

These data indicate a major role of CBS in intracellular H<sub>2</sub>S biosynthesis compared to CTH in ASC52telo cells. In fact, *CBS* gene KD in ASC52telo cells resulted in a significant increase in cellular inflammation and oxidative stress in parallel to decreased capacity to produce endogenous H<sub>2</sub>S. A large number of studies demonstrated anti-inflammatory and anti-oxidant effects of H<sub>2</sub>S in mesenchymal stem

cells [10,12] and other cells [13–15]. However, the pro-oxidant and pro-inflammatory effects of CBS depletion might be also explained by other causes, including:

- Altered mitochondrial respiratory function, characterized by decreased proton leak and spare respiratory capacity, two functional measures of mitochondrial function [16,17]. Supporting these findings, a recent study demonstrated the importance of CBS on mitochondrial function in endothelial cells [18]. It is important to note that proton leak (uncoupled respiration), which was significantly decreased in CBS KD ASC52telo (shCBS-ASC52telo) cells (Fig. 2K), is key mitochondrial process in the prevention of oxidative stress [19,20].
- Decreased expression of rejuvenation (*LC3* and *SIRT1*)-related genes. The protective effects of increased *SIRT1* [21–27] and *LC3* [28–31] mRNA levels in the prevention of cellular inflammation, oxidative stress and mitochondrial function are widely reported.
- Increased homocysteine/cystathioneine ratio. CBS enzyme inhibition results in increased homocysteine in detrimental of cystathione levels [32,33]. The pro-oxidant and pro-inflammatory effects of excess homocysteine have been demonstrated in *in vitro* and *in vivo* experiments [33–38]. Otherwise, similar to ASC52telo cells, breast tumor tissue displayed increased CBS, but very low CTH mRNA levels, resulting in increased intracellular levels of cystathione, and in consequence preserving mitochondrial



**Fig. 5. A–J)** Effect of permanent CBS gene knockdown (shCBS) on *CBS* mRNA levels (A), expression of osteogenic-related genes (*RUNX2*, *BGLAP*, *TGFB1*) (B–D), rejuvenation-related genes (*LC3*, *SIRT1*) (E–F), *FABP4* (G) and proinflammatory cytokines (*IL6*, *TNF*, *CXCL8*) (H–J) during ASC52telo cell osteocyte differentiation (at day 9). \*p < 0.05, \*\*p < 0.01 and \*\*\*p < 0.001 vs shC-ASC52telo cells.

function and preventing oxidative and endoplasmic reticulum stress [39].

Another important finding of current study was that *CBS* gene knockdown in ASC52telo cells greatly enhanced their capacity to differentiate into adipocytes in detriment to the ability to differentiate into osteogenic lineage. It is well-established that increased oxidative stress (ROS levels) in adipose-derived mesenchymal stem cells promotes a cellular senescence and inflammation phenotype characterised by increased capacity for differentiate into adipocyte at the expense of decreased stemness capacity [40–42]. In fact, antioxidant treatment in mesenchymal stem cells prevented adipocyte differentiation [42]. In agreement with current data, the administration of diallyl disulfide, a slow H<sub>2</sub>S donor [43], restored hAMSC stemness via inhibition of intracellular ROS levels [44]. The relevance of *CBS* on osteogenesis has been previously reported [45–47]. Interestingly, one of these studies demonstrated that exogenous homocysteine administration increased intracellular homocysteine levels, and resulted in increased ROS, altered mitochondrial function, reduced expression of *CBS* gene and intracellular H<sub>2</sub>S level, and in consequence, inhibiting its capacity to differentiate into osteogenic lineage [47]. Otherwise, to the best of our knowledge, the impact of *CBS* depletion enhancing hAMSC adipogenesis has not been previously shown. These findings could seem controversial with one previous study in 3T3-L1 cells, in which *CBS* gene knockdown decreased adipogenesis [1], whereas other studies demonstrated that CTH was the only enzyme of transulfuration pathway with adipogenic properties [2–4]. In line with these studies, increased CTH/CBS ratio was associated to increased adipogenic potential in ASC52telo cells, which was characterised with very high expression of adipogenic genes (such as *FABP4* and *SLC2A4*), even before adding the adipogenic media

(at day 0). Moreover, supporting the cellular inflammation-associated adipogenic potential observed in shCBS-ASC52telo cells, two recent studies demonstrated that cellular inflammation promoted the generation of new adipocytes and adipose tissue expansion [48,49]. Otherwise, *IRS1* gene expression was significantly decreased in shCBS-ASC52telo cells at day 0 and during adipocyte differentiation. In line with this, even though IRS-1 is important for lipid storage in adipose tissue [50], a recent study reported that IRS-1 knockdown in bone marrow mesenchymal stem cells increased their capacity to differentiate into adipocytes [51]. Current findings and these studies [45,46,48] suggest that cellular senescence and inflammation might promote a non-physiological increased adipocyte differentiation.

## 5. Conclusions

Altogether these findings demonstrated that permanent *CBS* gene KD in ASC52telo cells promotes a cellular senescence phenotype, characterized by increased cellular inflammation and oxidative stress, reduced cellular rejuvenation-related gene expression markers and a very increased adipogenic potential, which resulted in a non-physiological enhanced adipocyte differentiation with excessive lipid storage.

## Author contributions

FC, JL, FO, NO-C and AL researched data; WR and JMF-R contributed to the discussion and reviewed the manuscript; JMMN researched data and wrote the manuscript.

## Declaration of competing interest

The authors have nothing to disclose.

## Acknowledgements

This work was partially supported by research grants PI15/01934, PI16/01173 and PI19/01712 from the Instituto de Salud Carlos III from Spain and VII Spanish Diabetes Association grants to Basic Diabetes Research Projects led by young researchers, CIBERON Fisiopatología de la Obesidad y Nutrición is an initiative from the Instituto de Salud Carlos III and Fondo Europeo de Desarrollo Regional (FEDER) from Spain. We acknowledge the technical support in L1 synthesis of Xavier Ribas (UdG) and Miquel Costas (UdG).

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.redox.2020.101668>.

## References

- [1] C.Y. Tsai, M.T. Peh, W. Feng, B.W. Dymock, P.K. Moore, Hydrogen sulfide promotes adipogenesis in 3T3L1 cells, *PLoS One* 10 (2015) e0119511–e0119511.
- [2] G. Yang, Y. Ju, M. Fu, Y. Zhang, Y. Pei, M. Racine, S. Bath, T.J.S. Merritt, R. Wang, L. Wu, Cystathione gamma-lyase/hydrogen sulfide system is essential for adipogenesis and fat mass accumulation in mice, *Biochim. Biophys. Acta Mol. Cell Biol. Lipids* 1863 (2018) 165–176.
- [3] J. Cai, X. Shi, H. Wang, J. Fan, Y. Feng, X. Lin, J. Yang, Q. Cui, C. Tang, G. Xu, B. Geng, Cystathione  $\gamma$  lyase-hydrogen sulfide increases peroxisome proliferator-activated receptor  $\gamma$  activity by sulphydratation at C139 site thereby promoting glucose uptake and lipid storage in adipocytes, *Biochim. Biophys. Acta Mol. Cell Biol. Lipids* 1861 (2016) 419–429.
- [4] F. Comas, J. Latorre, O. Cussó, F. Ortega, A. Lluch, M. Sabater, A. Castells-Nobau, W. Ricart, X. Ribas, M. Costas, J.M. Fernández-Real, J.M. Moreno-Navarrete, Hydrogen sulfide impacts on inflammation-induced adipocyte dysfunction, *Food Chem. Toxicol.* 131 (2019) 110543.
- [5] Z. Pan, Z. Zhou, H. Zhang, H. Zhao, P. Song, D. Wang, J. Yin, W. Zhao, Z. Xie, F. Wang, Y. Li, C. Guo, F. Zhu, L. Zhang, Q. Wang, CD90 serves as differential modulator of subcutaneous and visceral adipose-derived stem cells by regulating AKT activation that influences adipose tissue and metabolic homeostasis, *Stem Cell Res. Ther.* 10 (2019).
- [6] B. Gustafson, A. Nerstedt, U. Smith, Reduced subcutaneous adipogenesis in human hypertrophic obesity is linked to senescent precursor cells, *Nat. Commun.* 10 (2019).
- [7] S. Wolbank, G. Stadler, A. Peterbauer, A. Gillich, M. Karbienier, B. Streubel, M. Wieser, H. Katinger, M. Van Grienden, H. Redl, C. Gabriel, J. Grillari, R. Grillari-Voglauer, Telomerase immortalized human amnion- and adipose-derived mesenchymal stem cells: maintenance of differentiation and immunomodulatory characteristics, *Tissue Eng. A* 15 (2009) 1843–1854.
- [8] S.A. Choi, C.S. Park, O.S. Kwon, H.K. Giong, J.S. Lee, T.H. Ha, C.S. Lee, Structural effects of naphthalimide-based fluorescent sensor for hydrogen sulfide and imaging in live zebrafish, *Sci. Rep.* 6 (2016), 26203–26203.
- [9] J.M. Moreno-Navarrete, M. Jové, T. Padró, J. Boada, F. Ortega, W. Ricart, R. Pamplona, L. Badimon, M. Portero-Otín, J.M. Fernández-Real, Adipocyte lipopolysaccharide binding protein (LBP) is linked to a specific lipidomic signature, *Obesity* 25 (2017) 391–400.
- [10] B. Fox, J.T. Schantz, R. Haigh, M.E. Wood, P.K. Moore, N. Viner, J.P.E. Spencer, P. G. Winyard, M. Whiteman, Inducible hydrogen sulfide synthesis in chondrocytes and mesenchymal progenitor cells: is H2S a novel cytoprotective mediator in the inflamed joint? *J. Cell Mol. Med.* 16 (2012) 896–910.
- [11] T.A. Markel, N.A. Drucker, A.R. Jensen, K.R. Olson, Human mesenchymal stem cell hydrogen sulfide production critically impacts the release of other paracrine mediators after injury, *J. Surg. Res.* 254 (2020) 75–82.
- [12] I. Cacciotti, M. Ciocci, E. Di Giovanni, F. Nanni, S. Melino, Hydrogen sulfide-releasing fibrous membranes: potential patches for stimulating human stem cells proliferation and viability under oxidative stress, *Int. J. Mol. Sci.* 19 (2018).
- [13] J. Li, X. Teng, S. Jin, J. Dong, Q. Guo, D. Tian, Y. Wu, Hydrogen sulfide improves endothelial dysfunction by inhibiting the vicious cycle of NLRP3 inflammasome and oxidative stress in spontaneously hypertensive rats, *J. Hypertens.* 37 (2019) 1633–1643.
- [14] Z. Lin, N. Altaf, C. Li, M. Chen, L. Pan, D. Wang, L. Xie, Y. Zheng, H. Fu, Y. Han, Y. Ji, Hydrogen sulfide attenuates oxidative stress-induced NLRP3 inflammasome activation via S-sulphydratation of c-Jun at Cys269 in macrophages, *Biochim. Biophys. Acta (BBA) - Mol. Basis Dis.* 1864 (2018) 2890–2900.
- [15] W. P, C. F, W. W, Z. XD, Hydrogen sulfide attenuates high glucose-induced human retinal pigment epithelial cell inflammation by inhibiting ROS formation and NLRP3 inflammasome activation, *Mediat. Inflamm.* 2019 (2019).
- [16] Y.X. Xiao, I.R. Lanza, J.M. Swain, M.G. Sarr, K.S. Nair, M.D. Jensen, Adipocyte mitochondrial function is reduced in human obesity independent of fat cell size, *J. Clin. Endocrinol. Metab.* 99 (2014).
- [17] J.M. Moreno-Navarrete, F. Ortega, A. Rodríguez, J. Latorre, S. Becerril, M. Sabater-Masdeu, W. Ricart, G. Frühbeck, J.M. Fernández-Real, HMOX1 as a marker of iron excess-induced adipose tissue dysfunction, affecting glucose uptake and respiratory capacity in human adipocytes, *Diabetologia* 60 (2017) 915–926.
- [18] G. Rao, B. Murphy, A. Dey, S.K.D. Dwivedi, Y. Zhang, R.V. Roy, P. Chakraborty, R. Bhattacharya, P. Mukherjee, Cystathione beta synthase regulates mitochondrial dynamics and function in endothelial cells, *Faseb. J.* (2020), <https://doi.org/10.1096/fj.202000173R>.
- [19] S. Cadena, Mitochondrial uncoupling, ROS generation and cardioprotection, *Biochim. Biophys. Acta Bioenerg.* 1859 (2018) 940–950.
- [20] P.S. Brookes, Mitochondrial H<sup>+</sup> leak and ROS generation: an odd couple, *Free Radic. Biol. Med.* 38 (2005) 12–23.
- [21] G.-Z. Zhang, Y.-J. Deng, Q.-Q. Xie, E.-H. Ren, Z.-J. Ma, X.-G. He, Y.-C. Gao, X.-W. Kang, Sirtuins and intervertebral disc degeneration: roles in inflammation, oxidative stress, and mitochondrial function, *Clin. Chim. Acta* 508 (2020) 33–42.
- [22] L. S., B. SH, K. YJ, P. JH, D.L. B, J. WB, J. S., L. NK, D. L, K. DY, K. S., S. HJ, Y. J, L. DH, M. HR, C. HY, K. SM, MHY2233 attenuates replicative cellular senescence in human endothelial progenitor cells via SIRT1 signaling, *Oxid. Med. Cell. Longev.* 2019 (2019), 6492029–6492029.
- [23] J.R. Baker, C. Vuppulsetty, T. Colley, S. Hassibi, P.S. Fenwick, L.E. Donnelly, K. Ito, P.J. Barnes, MicroRNA-570 is a novel regulator of cellular senescence and inflammaging, *Faseb. J.* 33 (2019) 1605–1616.
- [24] P. Wang, C. Lv, T. Zhang, J. Liu, J. Yang, F. Guan, T. Hong, FOXQ1 regulates senescence-associated inflammation via activation of SIRT1 expression, *Cell Death Dis.* 8 (2017), e2946.
- [25] Y. Guo, P. Li, L. Gao, J. Zhang, Z. Yang, G. Bledsoe, E. Chang, L. Chao, J. Chao, Kallistatin reduces vascular senescence and aging by regulating microRNA-34a-SIRT1 pathway, *Aging Cell* 16 (2017) 837–846.
- [26] Y. Yuan, V.F. Cruzat, P. Newsholme, J. Cheng, Y. Chen, Y. Lu, Regulation of SIRT1 in aging: roles in mitochondrial function and biogenesis, *Mech. Ageing Dev.* 155 (2016) 10–21.
- [27] J.W. Hwang, H. Yao, S. Caito, I.K. Sundar, I. Rahman, Redox regulation of SIRT1 in inflammation and cellular senescence, *Free Radic. Biol. Med.* 61 (2013) 95–110.
- [28] S. D'Adamo, S. Cetrullo, S. Guidotti, Y. Silvestri, M. Minguzzi, S. Santi, L. Cattini, G. Filardo, F. Flamigni, R.M. Borzi, Spermidine rescues the deregulated autophagic response to oxidative stress of osteoarthritic chondrocytes, *Free Radic. Biol. Med.* 153 (2020) 159–172.
- [29] J. W., Z. P., Z. X. Apelin promotes ECM synthesis by enhancing autophagy Flux via TFEB in human degenerative NP cells under oxidative stress, *BioMed Res. Int.* 2020 (2020), 4897170–4897170.
- [30] S. Bhansali, A. Bhansali, P. Dutta, R. Walia, V. Dhawan, Metformin upregulates mitophagy in patients with T2DM: a randomized placebo-controlled study, *J. Cell Mol. Med.* 24 (2020) 2832–2846.
- [31] R. Li, J. Shang, W. Zhou, L. Jiang, D. Xie, G. Tu, Overexpression of HIPK2 attenuates spinal cord injury in rats by modulating apoptosis, oxidative stress, and inflammation, *Biomol. Pharmacother.* 103 (2018) 127–134.
- [32] K. Zuhra, F. Augsburger, T. Majtan, C. Szabo, Cystathione- $\beta$ -synthase: molecular regulation and pharmacological inhibition, *Biomolecules* 10 (2020).
- [33] K. Robert, J. Nehmé, E. Bourdon, G. Pivert, B. Friguet, C. Delcayre, J. Delabar, N. Janel, Cystathione  $\beta$  synthase deficiency promotes oxidative stress, fibrosis, and steatosis in mice liver, *Gastroenterology* 128 (2005) 1405–1415.
- [34] K. AnandBabu, P. Sen, N. Angayarkanni, Oxidized LDL, homocysteine, homocysteine thiolactone and advanced glycation end products act as pro-oxidant metabolites inducing cytokine release, macrophage infiltration and pro-angiogenic effect in ARPE-19 cells, *PLoS One* 14 (2019) e0216899–e0216899.
- [35] X. Wu, L. Zhang, Y. Miao, J. Yang, X. Wang, C. Chen Wang, J. Feng, L. Wang, Homocysteine causes vascular endothelial dysfunction by disrupting endoplasmic reticulum redox homeostasis, *Redox Biol.* 20 (2019) 46–59.
- [36] A. Longoni, B. Bellaver, L.D. Bobermin, C.L. Santos, Y. Nonose, J. Kolling, T.M. dos Santos, A.M. de Assis, A. Quincozes-Santos, A.T.S. Wyse, Homocysteine induces glial reactivity in adult rat astrocyte cultures, *Mol. Neurobiol.* 55 (2018) 1966–1976.
- [37] D. Zhang, P. Fang, X. Jiang, J. Nelson, J.K. Moore, W.D. Kruger, R.M. Berretta, S. R. Houser, X. Yang, H. Wang, Severe hyperhomocysteineemia promotes bone marrow-derived and resident inflammatory monocyte differentiation and atherosclerosis in LDLr/CBS-deficient mice, *Circ. Res.* 111 (2012) 37–49.
- [38] L. Papatheodorou, N. Weiss, Vascular oxidant stress and inflammation in hyperhomocysteinemia, *Antioxidants Redox Signal.* 9 (2007) 1941–1958.
- [39] S. Sen, B. Kawahara, S.K. Mahata, R. Tsai, A. Yoon, L. Hwang, K. Hu-Moore, C. Villanueva, A. Vajihuddin, P. Parameshwar, M. You, D.L. Bhaskar, O. Gomez, K. F. Faull, R. Farias-Eisner, G. Chaudhuri, Cystathione: a novel oncometabolite in human breast cancer, *Arch. Biochem. Biophys.* 604 (2016) 95–102.
- [40] C.H. Lin, N.T. Li, H.S. Cheng, M.L. Yen, Oxidative stress induces imbalance of adipogenic/osteoblastic lineage commitment in mesenchymal stem cells through decreasing SIRT1 functions, *J. Cell Mol. Med.* 22 (2018) 786–796.
- [41] K. Marycz, K.A. Tomaszewski, K. Kornicka, B.M. Henry, S. Wroński, J. Tarasiuk, M. Maredzianik, Metformin decreases reactive oxygen species, enhances osteogenic properties of adipose-derived multipotent mesenchymal stem cells in vitro, and increases bone density in vivo, *Oxid. Med. Cell. Longev.* 2016 (2016) 9785890.
- [42] D. de Villiers, M. Potgieter, M.A. Ambele, L. Adam, C. Durandt, M.S. Pepper, The role of reactive oxygen species in adipogenic differentiation, in: *Adv. Exp. Med. Biol.*, Springer New York LLC, 2018, pp. 125–144.

- [43] D. Liang, H. Wu, M.W. Wong, D. Huang, Diallyl trisulfide is a fast H<sub>2</sub>S donor, but diallyl disulfide is a slow one: the reaction pathways and intermediates of glutathione with polysulfides, *Org. Lett.* 17 (2015) 4196–4199.
- [44] K. Baharmpour Juybari, T. Kamarul, M. Najafi, D. Jafari, A.M. Sharifi, Restoring the IL-1 $\beta$ /NF- $\kappa$ B-induced impaired chondrogenesis by diallyl disulfide in human adipose-derived mesenchymal stem cells via attenuation of reactive oxygen species and elevation of antioxidant enzymes, *Cell Tissue Res.* 373 (2018) 407–419.
- [45] Y. Liu, R. Yang, X. Liu, Y. Zhou, C. Qu, T. Kikuri, S. Wang, E. Zandi, J. Du, I. S. Ambudkar, S. Shi, Hydrogen sulfide maintains mesenchymal stem cell function and bone homeostasis via regulation of Ca<sup>2+</sup> channel sulfhydration, *Cell Stem Cell* 15 (2014) 66–78.
- [46] L. Gambari, G. Lisignoli, E. Gabusi, C. Manferdini, F. Paoletta, A. Piacentini, F. Grassi, Distinctive expression pattern of cystathionine- $\beta$ -synthase and cystathionine- $\gamma$ -lyase identifies mesenchymal stromal cells transition to mineralizing osteoblasts, *J. Cell. Physiol.* 232 (2017) 3574–3585.
- [47] Y. Zhai, J. Behera, S.C. Tyagi, N. Tyagi, Hydrogen sulfide attenuates homocysteine-induced osteoblast dysfunction by inhibiting mitochondrial toxicity, *J. Cell. Physiol.* 234 (2019) 18602–18614.
- [48] Q. Zhu, Y.A. An, M. Kim, Z. Zhang, S. Zhao, Y. Zhu, J.W. Asterholm, C. M. Kusminski, P.E. Scherer, Suppressing adipocyte inflammation promotes insulin resistance in mice, *Mol. Metab.* 39 (2020) 101010.
- [49] I. Wernstedt Asterholm, C. Tao, T.S. Morley, Q.A. Wang, F. Delgado-Lopez, Z. V. Wang, P.E. Scherer, Adipocyte inflammation is essential for healthy adipose tissue expansion and remodeling, *Cell Metabol.* 20 (2014) 103–118.
- [50] Y.H. Tseng, A.J. Butte, E. Kokkotou, V.K. Yechoor, C.M. Taniguchi, K. M. Kriauciunas, A.M. Cypess, M. Niimobe, K. Yoshikawa, M.E. Patti, C.R. Kahn, Prediction of preadipocyte differentiation by gene expression reveals role of insulin receptor substrates and necdin, *Nat. Cell Biol.* 7 (2005) 601–611.
- [51] N. Wang, Y. Li, Z. Li, J. Ma, X. Wu, R. Pan, Y. Wang, L. Gao, X. Bao, P. Xue, IRS-1 targets TAZ to inhibit adipogenesis of rat bone marrow mesenchymal stem cells through PI3K-Akt and MEK-ERK pathways, *Eur. J. Pharmacol.* 849 (2019) 11–21.