



INORGANIC CHEMISTRY

FRONTIERS

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: C. Magallon, L. Griego, C. Hu, A. Company, X. Ribas and L. M. Mirica, *Inorg. Chem. Front.*, 2022, DOI: 10.1039/D1QI01486B.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Organometallic Ni(II), Ni(III), and Ni(IV) Complexes Relevant to Carbon-Carbon and Carbon-Oxygen Bond Formation Reactions

C. Magallon,^a L. Griego,^b C. H. Hu,^b A. Company,^{*a} X. Ribas^{*a} and L. M. Mirica^{*b}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Herein, the pyridinophane tetradentate ligand 3,6,9-trimethyl-3,6,9,15-Tetraazabicyclo[9.3.1]pentadeca-1(15),11,13-triene, PyNMe₃, is used to isolate and structurally characterize well-defined organometallic Ni(II) and Ni(III) complexes bearing the cycloneophyl fragment, an alkyl/aryl C-donor ligand. Furthermore, spectroscopic and cryo-mass spectrometry studies suggest the formation of a transient Ni(IV) organometallic complex, and its relevance to C-C and C-O bond formation reactivity studies is discussed.

Introduction

The formation of new C-C and C-heteroatom bonds through transition-metal catalyzed cross-coupling reactions (i.e., Suzuki, Kumada, and Negishi reactions) is currently one of the most powerful tools in organic chemistry.¹⁻³ In this context, noble metal-based systems are the more commonly used catalysts for these transformations, due to their widely known mechanistic understanding.⁴⁻⁷ For example, Pd-catalyzed reactions usually occur via Pd⁰/Pd^{II} catalytic cycles and mostly involve diamagnetic intermediate species, which makes their characterization more feasible.⁸ In contrast, the mechanisms of Ni-catalyzed cross-coupling reactions are far less understood since this first-row transition-metal can easily undergo both one- and two-electron redox processes, often involving paramagnetic intermediate species that lead to more complex mechanistic pathways.⁹⁻¹¹ Although traditionally Ni-catalyzed cross-coupling reactions have been proposed to involve Ni⁰, Ni^I and Ni^{II} intermediates,¹²⁻¹⁴ recent studies suggest that high-valent Ni^{III} and Ni^{IV} species are key intermediates in the C-C/C-heteroatom bond-forming step.¹⁵⁻²⁴

Pyridinophane tetradentate ligands have been shown to stabilize uncommon high-valent organometallic nickel complexes. Indeed, in the past several years, Mirica and co-workers have employed a series of tetradentate pyridinophane ligands with different amine N-substituents (^RN₄ ligands, R = Me, Ts, ^tPr, ^tBu, Np) as well as the tridentate 1,4,7-trimethyl-1,4,7-

triazacyclononane ligand (Me₃tacn) to stabilize Ni^{III/IV} complexes (Fig. 1a), as well as Pd^{II/IV} complexes, that undergo C-C/C-heteroatom bond formation reactions.²⁵⁻⁴⁰ Recently, a macrocyclic N-based tetradentate ligand PyNMe₃ has been employed by Costas, Company and co-workers to stabilize high-valent intermediate species and has proved successful to trap Fe^{IV/V}-oxo species (Fig. 1b) and dinuclear Cu^{II} side-on peroxy species.⁴¹⁻⁴⁵ However, the use of the PyNMe₃ ligand has




Fig. 1 a) Isolated organometallic nickel complexes bearing tridentate or tetradentate N-donor ligands and cycloneophyl as the C-donor ligand; b) Spectroscopically characterized high-valent iron-oxo intermediate species relevant for O₂ activation chemistry bearing the tetradentate N-based PyNMe₃ ligand, and c) Synthesis of well-defined organometallic nickel complexes bearing the PyNMe₃ and the cycloneophyl ligands and their C-C and C-O bond formation reactivity.

^a Institut de Química Computacional i Catàlisi (IQCC) and Departament de Química, Universitat de Girona, Campus Montilivi, Girona, E-17003, Catalonia, Spain.

^b Department of Chemistry, University of Illinois at Urbana-Champaign, 600 S. Mathews Avenue, Urbana, Illinois 61801, USA.

*Corresponding authors: anna.company@udg.edu; xavi.ribas@udg.edu; mirica@illinois.edu

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

never been employed to explore the reactivity of organometallic Ni complexes.

Herein we report the synthesis, characterization, and initial reactivity studies of a series of organometallic Ni^{II}, Ni^{III} and Ni^{IV} complexes supported by the PyNMe₃ ligand and containing the cycloneophyl fragment, an alkyl/aryl C-donor ligand (Fig. 1c). The cycloneophyl ligand was developed by Carmona *et al.* as an ancillary ligand to isolate organonickel complexes, since the resulting Ni(cycloneophyl) complexes are less prone to reductive elimination or β-hydride elimination.⁴⁶⁻⁴⁸ In addition, the cycloneophyl moiety has been widely and successfully employed together with N-based ligands independently by Sanford and Mirica in order to study C-C and C-heteroatom bond forming reactions.^{31-33, 49-54} The current study reports the reactivity of the isolated (PyNMe₃)Ni^{II}(cycl) complex with oxidants to promote C-C or C-O reductive elimination through Ni^{IV} intermediate species is described.

Results and Discussion

Our first attempt to isolate organometallic nickel complexes bearing the PyNMe₃ ligand consisted of the *in situ* oxidative addition of 1-bromo-4-fluorobenzene using Ni(cod)₂. The putative [(PyNMe₃)Ni^{II}(pF-Ph)(Br)] complex could not be isolated due to fast decomposition and rapid ligand exchange that leads to the formation of 0.5 equiv (PyNMe₃)Ni^{II}Br₂ (**1-Br**) and 0.5 equiv [(PyNMe₃)Ni^{II}(pF-Ph)₂]. The latter di-aryl complex then undergoes reductive elimination to afford the homocoupled product pF-Ph-Ph-pF (as detected via GC-MS), free ligand and Ni(0) (Fig. 2a). In addition, reaction of [(PyNMe₃)Ni^{II}(Cl)₂] (**1-Cl**) with MeMgCl in THF at -50 °C led to an intractable mixture due to fast decomposition of the desired nickel(II)-dimethyl complex (Fig. 2b). Nonetheless, the putative [(PyNMe₃)Ni^{II}(Me)₂] species could only be detected via ¹H NMR, which revealed two singlets below 0 ppm that integrated to three protons each and presumably correspond to the two inequivalent methyl groups directly attached to the Ni center (Figure S1).




Fig. 2 a) Attempt to synthesize [(PyNMe₃)Ni^{II}(PhF)(Br)] via oxidative addition at nickel(0), and b) Attempt to synthesize [(PyNMe₃)Ni^{II}(Me)₂] via transmetalation at the nickel(II)-chloride precursor, **1-Cl**.

2 | *J. Name.*, 2012, **00**, 1-3

Gratifyingly, the combination of PyNMe₃ and cycloneophyl (-CH₂CMe₂-o-C₆H₄-, or cycl) as ligands allows the stabilization and isolation of complex [(PyNMe₃)Ni^{II}(cycl)] (**2**) in 67 % yield. The complex was synthesized via ligand exchange of the Ni(II) precursor (py)₂Ni^{II}(cycl) with an equimolar amount of PyNMe₃ in 1:1 toluene/pentane, for 16 hours at room temperature under a nitrogen atmosphere (Fig. 3a). The X-ray structure revealed a square planar geometry around the Ni(II) center, with the PyNMe₃ ligand coordinated in a bidentate fashion (Fig. 3b). This new system allows for the observation the *trans* influence of the ligands. Interestingly, PyNMe₃ ligand coordinates to the Ni(II) center through the pyridine moiety which is *trans* to the alkyl ligand, and one of the adjacent amines *trans* to the phenyl group. This type of coordination could not be observed with the previously reported systems bearing symmetric pyridinophane type ligands (^RN4). In addition, the ¹H-NMR spectrum confirms that **2** is a low-spin diamagnetic Ni(II) complex. Detailed NMR experiments were used to assign all proton and carbon peaks for **2** (Figures S2-S9). The broadness of the peaks, especially the methylene protons on the PyNMe₃ ligand, is likely due to rapid ligand exchange of the four types of N donors. Furthermore, the unusual Ni(II) structure allows for an anagostic interaction of the methylene protons vicinal to the tertiary amine and the Ni metal center, with a Ni···H distance of 2.40 Å and a Ni···H-C9 angle of 170°. The cyclic voltammogram (CV) of **2** in 0.1 M (nBu₄N)PF₆/MeCN reveals a quasi-reversible redox event at E_{1/2} = 250 mV vs ferrocene (Fc), followed by another quasi-reversible oxidation at E_{1/2} = 630 mV (Fig. 3c). These observed pseudo-reversible oxidations are presumably assigned to the Ni^{III/II} and Ni^{IV/III} redox couples, respectively. This result suggests accessible Ni(III) and Ni(IV) species, likely due to the stabilizing effect of the two anionic chelating C donors from the cycloneophyl ligand, as well as the tetradentate PyNMe₃ ligand, as reported previously for high-valent iron-oxo species.^{27, 33, 37}




Fig. 3 a) Synthesis and characterization of complex [(PyNMe₃)Ni^{II}(cycl)], **2**; b) ORTEP representation of **2** (50% probability thermal ellipsoids; hydrogen atoms have been omitted for clarity; selected distances (Å): Ni-C24, 1.888(2); Ni-C15, 1.932(2); Ni-N2, 2.0535(19); Ni-N1, 1.9923(19), and c) CV of **2** in 0.1 M NBu₄(PF₆)/MeCN at a scan rate of 100 mV/s (Ag wire as reference electrode).

This journal is © The Royal Society of Chemistry 20xx

In fact, **2** was rapidly oxidized with 1 equiv of ferrocenium hexafluorophosphate (FcPF_6) in MeCN at -35°C to yield the reddish product $[(\text{PyNMe}_3)\text{Ni}^{\text{III}}(\text{cycl})](\text{PF}_6)$, **3** (Fig. 4a). The EPR spectrum of **3** in 1:3 MeCN/PrCN is consistent with a Ni(III) d^7 complex and exhibits a pseudoaxial signal with superhyperfine coupling to two nitrogen atoms in the g_z direction ($A_{2\text{N}} = 13.7\text{ G}$), indicating that two $I = 1$ nitrogen atoms, presumably from the ligand and/or solvent, coordinate to the Ni(III) center (Fig. 4b). Luckily, complex **3** was stable and isolated at low temperature in a 73 % yield.




Fig. 4 a) Synthesis and characterization of complex $[(\text{PyNMe}_3)\text{Ni}^{\text{III}}(\text{cycl})](\text{PF}_6)$, **3**; b) EPR spectra of complex **3** at 77 K 1:3 MeCN/PrCN (black: experimental; red: simulated); c) ORTEP representation of **3** (50% probability thermal ellipsoids; hydrogen atoms have been omitted for clarity; selected distances (\AA): Ni-C1, 1.9492 (12); Ni-C2, 1.9693 (14); Ni-N1, 1.9944 (11); Ni-N2, 2.2543 (13); Ni-N3, 2.1463 (12); Ni-N4, 2.2470 (12), and d) Cryo-HR-MS of **3** showing the monocationic peak (3^+) at $m/z = 438.2289$ (top) and simulation of its isotopic pattern (bottom).

The single crystal X-ray structure of **3** confirmed the six-coordinate Ni(III) center in a distorted octahedral geometry where the 6 coordination positions are occupied by the PyNMe_3 and cycloneophyl ligands, thus confirming the EPR data (Fig. 4c). Intriguingly, both the Ni-N and Ni-C distances in **3** are longer than those in **2**, as previously observed for similar complexes, and likely due to the change in the coordination number from 4 to 6 upon oxidation.^{33, 37} Moreover, cryo-ESI-mass spectrometry at -40°C was performed to further characterize this Ni(III) complex. The MS spectrum showed a major monocationic peak with a $m/z = 438.2289$ and an isotopic pattern fully consistent with the calculated values for 3^+ , $[(\text{C}_{14}\text{H}_{24}\text{N}_4)\text{Ni}^{\text{III}}(\text{C}_{10}\text{H}_{12})]^+$ (Fig. 4d).

Attempts to synthesize the corresponding Ni(IV) complex were conducted through a stepwise oxidation of **2**. The oxidation of **2** to obtain **3** was monitored by low-temperature UV/vis spectroscopy (Fig. 5a and b). This experiment was performed at -40°C using a 0.5 mM solution of **2** in MeCN. An initial spectrum was recorded for the starting Ni(II) complex, which exhibited an absorption band with $\lambda_{\text{max}} = 462\text{ nm}$. For the one-electron oxidation of each complex monitored by low-temperature UV/vis, $\text{NO}(\text{SbF}_6)$ was used as oxidant in order to obtain a cleaner spectrum, since Fc^+ exhibits two absorption bands at 325 nm and 440 nm that could interfere with the detection of new species. Thus, addition of 1 equiv of NO^+ to **2** immediately yields **3**, which exhibits an absorption band with $\lambda_{\text{max}} = 513\text{ nm}$. Afterwards, another equivalent of the same oxidant was added to generate the new Ni(IV) complex, **4**. Upon addition of the second equivalent of oxidant the UV/vis spectrum instantly changed and showed the formation of two absorption bands with λ_{max} of 384 nm and 470 nm, and a shoulder at 620 nm (Fig. 5b). Subsequent injection of an aliquot into cryo-ESI-mass spectrometer confirmed the generation of species **4** with a dicationic major peak with a mass value of $m/z = 219.1120$ and an isotopic pattern fully consistent with the calculated value for $[(\text{C}_{14}\text{H}_{24}\text{N}_4)\text{Ni}^{\text{IV}}(\text{C}_{10}\text{H}_{12})]^{2+}$ (Fig. 5c). This novel complex was further characterized by NMR spectroscopy, where a stepwise injection of 2 equiv NO^+ resulted in the observation of a new Ni(IV) species (Figures S10-S11).




Fig. 5 a) Step-wise one-electron oxidation to nickel(IV) from **2** using $\text{NO}(\text{SbF}_6)$ as oxidant; b) UV-vis characterization of complexes **2**, **3** and **4** obtained by consecutive one-electron oxidation of **2** with $\text{NO}(\text{SbF}_6)$, respectively, and c) Cryo-HR-MS of **4** showing the dicationic peak (4^{2+}) at $m/z = 219.1120$ (top) and simulation of its isotopic pattern (bottom).

In line with the UV-vis results, monitoring the oxidation by NMR provided complementary information. A color change from orange to red-pink was observed upon the addition of 1 equiv of oxidant in CD_3CN at -35°C , while the obtained paramagnetic NMR spectrum supports the formation of a Ni(III) species. To no surprise, a second equivalent of oxidant allowed the formation of a new diamagnetic species, tentatively assigned as a Ni(IV) complex. The inherent broadness of the NMR spectrum of the putative Ni(IV) species is likely caused by the residual Ni(III), which made peak assignment difficult. However, the obtained Ni(IV) spectrum closely resembles those of the recently reported Ni^{IV}-cyclonophyl complexes supported by the *N*-based macrocycles pyridinophane (^{Me}N4) and triazacyclononane (^{Me}tacn).^{32, 37} Due to the highly electrophilic nature of the Ni(IV) species, a key structural feature observed in both ^{Me}N4 and ^{Me}tacn Ni^{IV}-cyclonophyl systems is a large shift of the Ni^{IV}-CH₂- protons from ~ 2.10 ppm in their Ni^{II}-CH₂- counterparts to 5.33 ppm and 5.50 ppm, respectively. A similar shift was observed with **4**, in which the Ni^{IV}-CH₂- protons shifted to 5.29 ppm, similar to what was observed for (^{Me}tacn)Ni^{IV}(cycl).³²

Next, we focused on the reactivity of **3** and **4** in C-C or C-O bond formation reactions. First, we exposed the Ni^{III} species **3** to high temperatures (80°C) to evaluate its stability, and it was found that the compound decomposed to an intractable mixture, which contains $\sim 15\%$ of the protodemetalation product *t*-butylbenzene, yet no reductive elimination products were obtained. Then, since the Ni(IV), **4**, was metastable, we studied its reactivity by generating it *in situ* through oxidation of **2** with a variety of two-electron oxidants such as XeF₂, 1-fluoro-2,4,6-trimethylpyridinium triflate (NFTPT), and PhI(OAc)₂, yet no appreciable amount of any reductive elimination products were observed. On the contrary, promising results were obtained

when exploring its reactivity with green oxidants such as O₂, H₂O₂, and ^tBuOOH (Fig. 6).

In these reactions, only trace amounts ($<1\%$) of protodemetalation product *t*-butylbenzene, **B**, was observed in GC-MS, indicating a nearly full conversion of **2** into corresponding Ni^{III}/Ni^{IV} species. The reaction of **2** with 2 equiv H₂O₂ in 2-MeTHF results in the formation of 11% **C** and 9% **A** as the primary organic products. The reaction of **2** with ^tBuOOH shows similar distribution of products. On the other hand, the reaction of **2** with dioxygen generated up to 59% of C-C coupled product **A**, exhibiting an appreciable C-C bond formation reactivity, which is proposed to occur via the intermediacy of the Ni(IV) species **4**. It is important to note that this tetradentate PyNMe₃ ligand system displays a different reactivity from the structurally related pyridinophane ^{RN}4 system,³³ where C-O reductive elimination was greatly enhanced. In that case, it is likely that a Ni^{IV}-hydroxo transient, yet metastable intermediate was generated and led to competing C-C/C-O reductive elimination steps, which has been observed for M^{IV} complexes (M = Pt, Pd) bearing ligands that are prone




Fig. 6 Reactivity studies of **2** in front different two-electron oxidants, and the quantification of the respective organic products by GC-MS.

to reductive elimination.^{32,33} By contrast, the [(PyNMe₃)Ni(cycl)] system seems to favor the C-C bond formation reactivity over the formation of any oxygen-containing products, possibly because the PyNMe₃ ligand is flexible enough to strongly bind to the Ni(IV) species and complete the octahedral coordination geometry, without allowing the coordination of any additional oxidant-derived, O-donor ligand.

Conclusion

In summary, we have been able to explore the rich organometallic redox chemistry of Ni by characterizing spectroscopically Ni^{II} (**2**), Ni^{III} (**3**) and Ni^{IV} (**4**) complexes supported by the tetradentate N-donor ligand PyNMe₃. **2** features a square-planar geometry with two pendant aliphatic amine moieties, which enter into the coordination sphere of the metal upon oxidation to a Ni(III), **3**, that features a distorted octahedral geometry. Further reaction of **3** with another equivalent of oxidant affords the putative Ni(IV), **4**. Reactivity studies indicate that the in situ formed Ni(IV) complex, [(PyNMe₃)Ni^{IV}(cycl)](SbF₆)₂ (**4**), promotes C-C coupling over C-O coupling despite the use of O-based oxidants, in contrast to the related pyridinophane ^RN₄ systems.³³ Overall, this new PyNMe₃ ligand system provides important insight in understanding at a molecular level the electronic properties of dissymmetric macrocyclic ligands on well-defined Ni^{II} (**2**), Ni^{III} (**3**) and Ni^{IV} (**4**) organometallic complexes and their role in C-C and C-O bond formation transformations.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We acknowledge financial support from NSF CHE-1925751 to L.M., MINECO-Spain for projects PID2019-104498GB-I00 to X.R., PID2019-106699GB-I00 to A.C. Generalitat de Catalunya is also acknowledged for project 2017SGR264. We thank UdG for a IFuG PhD grant to C.M. X.R. and A.C. are thankful for an ICREA Acadèmia award. The authors are grateful to STR-UdG for technical support.

Notes and references

‡ CCDC 2118410 (**1-Br**), 2118413 (**1-Cl**), 2118411 (**2**), 2118412 (**3**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif. The Supporting Information also provides additional information.

1. F. Diederich and A. de Meijere, *Metal-Catalyzed Cross-Coupling Reactions*, Wiley-VCH, Weinheim, Germany, 2nd edn., 2004.

- J. F. Hartwig, *Organotransition Metal Chemistry: From Bonding to Catalysis*, University Science Books, Sausalito, 2010.
- J. Terao and N. Kambe, Cross-Coupling Reaction of Alkyl Halides with Grignard Reagents Catalyzed by Ni, Pd, or Cu Complexes with π -Carbon Ligand(s), *Acc. Chem. Res.*, 2008, **41**, 1545-1554.
- J. Magano and J. R. Dunetz, Large-Scale Applications of Transition Metal-Catalyzed Couplings for the Synthesis of Pharmaceuticals, *Chem. Rev.*, 2011, **111**, 2177-2250.
- J.-P. Corbet and G. Mignani, Selected Patented Cross-Coupling Reaction Technologies, *Chem. Rev.*, 2006, **106**, 2651-2710.
- C. C. C. Johansson Seechurn, M. O. Kitching, T. J. Colacot and V. Snieckus, Palladium-Catalyzed Cross-Coupling: A Historical Contextual Perspective to the 2010 Nobel Prize, *Angew. Chem. Int. Ed.*, 2012, **51**, 5062-5085.
- J. F. Hartwig, Carbon-Heteroatom Bond-Forming Reductive Eliminations of Amines, Ethers, and Sulfides, *Acc. Chem. Res.*, 1998, **31**, 852-860.
- J. F. Hartwig, Carbon-heteroatom bond formation catalysed by organometallic complexes, *Nature*, 2008, **455**, 314-322.
- X. Hu, Nickel-catalyzed cross coupling of non-activated alkyl halides: a mechanistic perspective, *Chem. Sci.*, 2011, **2**, 1867-1886.
- V. B. Phapale and D. J. Cárdenas, Nickel-catalysed Negishi cross-coupling reactions: scope and mechanisms, *Chem. Soc. Rev.*, 2009, **38**, 1598-1607.
- T. T. Tsou and J. K. Kochi, Reductive coupling of organometals induced by oxidation. Detection of metastable paramagnetic intermediates, *J. Am. Chem. Soc.*, 1978, **100**, 1634-1635.
- T. T. Tsou and J. K. Kochi, Mechanism of oxidative addition. Reaction of nickel(0) complexes with aromatic halides, *J. Am. Chem. Soc.*, 1979, **101**, 6319-6332.
- J. Cornella, E. Gómez-Bengoia and R. Martín, Combined Experimental and Theoretical Study on the Reductive Cleavage of Inert C-O Bonds with Silanes: Ruling out a Classical Ni(0)/Ni(II) Catalytic Couple and Evidence for Ni(I) Intermediates, *J. Am. Chem. Soc.*, 2013, **135**, 1997-2009.
- C. A. Laskowski, D. J. Bungum, S. M. Baldwin, S. A. Del Ciello, V. M. Iluc and G. L. Hillhouse, Synthesis and Reactivity of Two-Coordinate Ni(I) Alkyl and Aryl Complexes, *J. Am. Chem. Soc.*, 2013, **135**, 18272-18275.
- L. M. Mirica, S. M. Smith and L. Griego, Organometallic Chemistry of High-Valent Ni(III) and Ni(IV) Complexes in *Nickel Catalysis in Organic Synthesis*, 2020, S. Ogoshi (Ed.) pp. 223-248.
- A. Joshi-Pangu, C.-Y. Wang and M. R. Biscoe, Nickel-Catalyzed Kumada Cross-Coupling Reactions of Tertiary Alkylmagnesium Halides and Aryl Bromides/Triflates, *J. Am. Chem. Soc.*, 2011, **133**, 8478-8481.
- A. S. Dudnik and G. C. Fu, Nickel-Catalyzed Coupling Reactions of Alkyl Electrophiles, Including Unactivated Tertiary Halides, To Generate Carbon-Boron Bonds, *J. Am. Chem. Soc.*, 2012, **134**, 10693-10697.
- N. D. Schley and G. C. Fu, Nickel-Catalyzed Negishi Arylations of Propargylic Bromides: A Mechanistic Investigation, *J. Am. Chem. Soc.*, 2014, **136**, 16588-16593.
- J. C. Tellis, D. N. Primer and G. A. Molander, Single-electron transmetalation in organoboron cross-coupling by photoredox/nickel dual catalysis, *Science*, 2014, **345**, 433-436.
- Z. Zuo, D. T. Ahneman, L. Chu, J. A. Terrett, A. G. Doyle and D. W. C. MacMillan, Merging photoredox with nickel catalysis: Coupling of α -carboxyl sp³-carbons with aryl halides, *Science*, 2014, **345**, 437-440.

21. J. Cornella, J. T. Edwards, T. Qin, S. Kawamura, J. Wang, C.-M. Pan, R. Gianatassio, M. Schmidt, M. D. Eastgate and P. S. Baran, Practical Ni-Catalyzed Aryl–Alkyl Cross-Coupling of Secondary Redox-Active Esters, *J. Am. Chem. Soc.*, 2016, **138**, 2174–2177.
22. S. Z. Tasker, E. A. Standley and T. F. Jamison, Recent advances in homogeneous nickel catalysis, *Nature*, 2014, **509**, 299–309.
23. J. R. Bour, N. M. Camasso, E. A. Meucci, J. W. Kampf, A. J. Canty and M. S. Sanford, Carbon–Carbon Bond-Forming Reductive Elimination from Isolated Nickel(III) Complexes, *J. Am. Chem. Soc.*, 2016, **138**, 16105–16111.
24. N. M. Camasso and M. S. Sanford, Design, synthesis, and carbon-heteroatom coupling reactions of organometallic nickel(IV) complexes, *Science*, 2015, **347**, 1218–1220.
25. J. R. Khusnutdinova, N. P. Rath and L. M. Mirica, Stable Mononuclear Organometallic Pd(III) Complexes and Their C–C Bond Formation Reactivity, *J. Am. Chem. Soc.*, 2010, **132**, 7303–7305.
26. J. R. Khusnutdinova, N. P. Rath and L. M. Mirica, Dinuclear Palladium(III) Complexes with a Single Unsupported Bridging Halide Ligand: Reversible Formation from Mononuclear Palladium(II) or Palladium(IV) Precursors, *Angew. Chem. Int. Ed.*, 2011, **50**, 5532–5536.
27. J. R. Khusnutdinova, N. P. Rath and L. M. Mirica, The Aerobic Oxidation of a Pd(II) Dimethyl Complex Leads to Selective Ethane Elimination from a Pd(III) Intermediate, *J. Am. Chem. Soc.*, 2012, **134**, 2414–2422.
28. F. Tang, Y. Zhang, N. P. Rath and L. M. Mirica, Detection of Pd(III) and Pd(IV) Intermediates during the Aerobic Oxidative C–C Bond Formation from a Pd(II) Dimethyl Complex, *Organometallics*, 2012, **31**, 6690–6696.
29. F. Tang, F. Qu, J. R. Khusnutdinova, N. P. Rath and L. M. Mirica, Structural and reactivity comparison of analogous organometallic Pd(III) and Pd(IV) complexes, *Dalton Trans.*, 2012, **41**, 14046–14050.
30. J. R. Khusnutdinova, N. P. Rath and L. M. Mirica, The Conformational Flexibility of the Tetradentate Ligand tBuN4 is Essential for the Stabilization of (tBuN4)Pd(III) Complexes, *Inorg. Chem.*, 2014, **53**, 13112–13129.
31. F. Qu, J. R. Khusnutdinova, N. P. Rath and L. M. Mirica, Dioxygen activation by an organometallic Pd(II) precursor: formation of a Pd(IV)–OH complex and its C–O bond formation reactivity, *Chem. Commun.*, 2014, **50**, 3036–3039.
32. M. B. Watson, N. P. Rath and L. M. Mirica, Oxidative C–C Bond Formation Reactivity of Organometallic Ni(II), Ni(III), and Ni(IV) Complexes, *J. Am. Chem. Soc.*, 2017, **139**, 35–38.
33. S. M. Smith, O. Planas, L. Gómez, N. P. Rath, X. Ribas and L. M. Mirica, Aerobic C–C and C–O bond formation reactions mediated by high-valent nickel species, *Chem. Sci.*, 2019, **10**, 10366–10372.
34. B. Zheng, F. Tang, J. Luo, J. W. Schultz, N. P. Rath and L. M. Mirica, Organometallic Nickel(III) Complexes Relevant to Cross-Coupling and Carbon–Heteroatom Bond Formation Reactions, *J. Am. Chem. Soc.*, 2014, **136**, 6499–6504.
35. F. Tang, N. P. Rath and L. M. Mirica, Stable bis(trifluoromethyl)nickel(III) complexes, *Chem. Commun.*, 2015, **51**, 3113–3116.
36. W. Zhou, J. W. Schultz, N. P. Rath and L. M. Mirica, Aromatic Methoxylation and Hydroxylation by Organometallic High-Valent Nickel Complexes, *J. Am. Chem. Soc.*, 2015, **137**, 7604–7607.
37. J. W. Schultz, K. Fuchigami, B. Zheng, N. P. Rath and L. M. Mirica, Isolated Organometallic Nickel(III) and Nickel(IV) Complexes Relevant to Carbon–Carbon Bond Formation Reactions, *J. Am. Chem. Soc.*, 2016, **138**, 12928–12934.
38. W. Zhou, S. Zheng, J. W. Schultz, N. P. Rath and L. M. Mirica, Aromatic Cyanoalkylation through Double C–H Activation Mediated by Ni(III), *J. Am. Chem. Soc.*, 2016, **138**, 5777–5780.
39. S. M. Smith, N. P. Rath and L. M. Mirica, Axial Donor Effects on Oxidatively Induced Ethane Formation from Nickel–Dimethyl Complexes, *Organometallics*, 2019, **38**, 3602–3609.
40. H. Na, M. B. Watson, F. Tang, N. P. Rath and L. M. Mirica, Photoreductive chlorine elimination from a Ni(III)Cl₂ complex supported by a tetradentate pyridinophane ligand, *Chem. Commun.*, 2021, **57**, 7264–7267.
41. E. Andris, J. Jašík, L. Gómez, M. Costas and J. Roithová, Spectroscopic Characterization and Reactivity of Triplet and Quintet Iron(IV) Oxo Complexes in the Gas Phase, *Angew. Chem. Int. Ed.*, 2016, **55**, 3637–3641.
42. J. Serrano-Plana, A. Aguinaco, R. Belda, E. García-España, M. G. Basallote, A. Company and M. Costas, Exceedingly Fast Oxygen Atom Transfer to Olefins via a Catalytically Competent Nonheme Iron Species, *Angew. Chem. Int. Ed.*, 2016, **55**, 6310–6314.
43. J. Serrano-Plana, W. N. Oloo, L. Acosta-Rueda, K. K. Meier, B. Verdejo, E. García-España, M. G. Basallote, E. Münck, L. Que, A. Company and M. Costas, Trapping a Highly Reactive Nonheme Iron Intermediate That Oxygenates Strong C–H Bonds with Stereoretention, *J. Am. Chem. Soc.*, 2015, **137**, 15833–15842.
44. V. Dantignana, J. Serrano-Plana, A. Draksharapu, C. Magallón, S. Banerjee, R. Fan, I. Gamba, Y. Guo, L. Que, M. Costas and A. Company, Spectroscopic and Reactivity Comparisons between Nonheme Oxoiron(IV) and Oxoiron(V) Species Bearing the Same Ancillary Ligand, *J. Am. Chem. Soc.*, 2019, **141**, 15078–15091.
45. C. Magallón, J. Serrano-Plana, S. Roldán-Gómez, X. Ribas, M. Costas and A. Company, Preparation of a coordinatively saturated μ - η^2 -peroxodicopper(II) compound, *Inorg. Chim. Acta*, 2018, **481**, 166–170.
46. E. Carmona, F. González, M. L. Poveda, J. L. Atwood and R. D. Rogers, Synthesis and properties of dialkyl complexes of nickel(II). The crystal structure of bis(pyridine)bis(trimethylsilylmethyl)nickel(II), *J. Chem. Soc., Dalton Trans.*, 1981, 777–782.
47. E. Carmona, E. Gutierrez-Puebla, J. M. Marin, A. Monge, M. Paneque, M. L. Poveda and C. Ruiz, Synthesis and x-ray structure of the nickelabenzocyclopentene complex [cyclic](Me₃P)₂Ni(CH₂CMe₂-o-C₆H₄). Reactivity toward simple, unsaturated molecules and the crystal and molecular structure of the cyclic carboxylate (Me₃P)₂Ni(CH₂CMe₂-o-C₆H₄C(O)O), *J. Am. Chem. Soc.*, 1989, **111**, 2883–2891.
48. E. Carmona, P. Palma, M. Paneque, M. L. Poveda, E. Gutierrez-Puebla and A. Monge, Synthesis of [cyclic]-(Me₃P)₂Ni(CH₂CMe₂-o-C₆H₄) and its reactivity toward carbon dioxide, carbon monoxide and formaldehyde. First observation of a carbonyl-carbonate oxidative conproportionation mediated by a transition-metal complex, *J. Am. Chem. Soc.*, 1986, **108**, 6424–6425.
49. N. M. Camasso, A. J. Canty, A. Ariafard and M. S. Sanford, Experimental and Computational Studies of High-Valent Nickel and Palladium Complexes, *Organometallics*, 2017, **36**, 4382–4393.
50. A. J. Canty, A. Ariafard, N. M. Camasso, A. T. Higgs, B. F. Yates and M. S. Sanford, Computational study of C(sp³)–O bond formation at a Pd(IV) centre, *Dalton Trans.*, 2017, **46**, 3742–3748.
51. I. M. Pendleton, M. H. Pérez-Temprano, M. S. Sanford and P. M. Zimmerman, Experimental and Computational Assessment of Reactivity and Mechanism in C(sp³)–N Bond-Forming

- Reductive Elimination from Palladium(IV), *J. Am. Chem. Soc.*, 2016, **138**, 6049-6060.
52. N. M. Camasso, M. H. Pérez-Temprano and M. S. Sanford, C(sp³)–O Bond-Forming Reductive Elimination from PdIV with Diverse Oxygen Nucleophiles, *J. Am. Chem. Soc.*, 2014, **136**, 12771-12775.
53. M. H. Pérez-Temprano, J. M. Racowski, J. W. Kampf and M. S. Sanford, Competition between sp³-C–N vs sp³-C–F Reductive Elimination from PdIV Complexes, *J. Am. Chem. Soc.*, 2014, **136**, 4097-4100.
54. J. M. Racowski, J. B. Gary and M. S. Sanford, Carbon(sp³)-Fluorine Bond-Forming Reductive Elimination from Palladium(IV) Complexes, *Angew. Chem. Int. Ed.*, 2012, **51**, 3414-3417.

View Article Online
DOI: 10.1039/D1QI01486B